

CHAPTER ONE: FUNDAMENTALS OF ALGEBRA**Review September 22** ↻ **Test September 29***

Thorough understanding and fluency of the concepts and methods in this chapter is a cornerstone to success in the rest of this course and most future math courses as well. The content will be covered very quickly since most of it is review, but nonetheless it is some of the most important content of the year (and of all mathematics).

1-A Fractions**Wednesday • 9/9**

- ① Convert a whole number to a fraction.
- ② Multiply a fraction.
- ③ Divide by a fraction.
- ④ Reduce a fraction.
- ⑤ Add or subtract fractions with the same denominator.
- ⑥ Add or subtract fractions with different denominators.
- ⑦ Convert a percentage to a decimal or fraction.

1-B Expressions**Friday • 9/11**

identity element • term • expression • argument • polynomial • monomial • binomial • trinomial • degree • constant • linear • quadratic • cubic • standard form • coefficient • leading coefficient • scientific notation

- ① Identify the terms of an expression and their coefficients.
- ② Identify the argument of a function.
- ③ Multiply a rational expression by an integer.
- ④ Simplify a fraction with multiple terms in the numerator or denominator.
- ⑤ Classify a polynomial in one variable.
- ⑥ Convert calculator notation to scientific notation and to standard notation.
- ⑦ Convert standard notation to scientific notation.

1-C Solving Equations**Tuesday • 9/15**

inverse • equation

- ① Apply order of operations.
- ② Identify the inverse of an operation by definition.
- ③ Solve an equation.
- ④ Show proper notation in solving an equation.
- ⑤ Express an answer.

1-D Properties of Exponents**Thursday • 9/17**

- ① Simplify an expression using properties of exponents.

1-E Addition, Subtraction, and Multiplication of Polynomials**Monday • 9/21**

conjugate

- ① Add or subtract polynomials.
- ② Multiply two polynomials.
- ③ Multiply more than two polynomials.
- ④ Multiply a binomial by its conjugate.
- ⑤ Square a binomial.

** This chapter's dates are subject to change due to the current unique circumstances.*

1-A Fractions

① Convert a whole number to a fraction.

1. The numerator is the original number, and the denominator is 1.

① 10 can be written as the fraction $\frac{10}{1}$.

② Multiply a fraction.

1. Multiply the numerators together.

2. Multiply the denominators together.

② a) $\frac{2}{5} \times \frac{3}{4} = \frac{6}{20}$ b) $\frac{2}{5} \times 3 = \frac{2}{5} \times \frac{3}{1} = \frac{6}{5}$

③ Divide by a fraction.

1. Multiply by the reciprocal of the fraction (see ②).

③ a) $\frac{2}{5} \div \frac{4}{3} = \frac{2}{5} \times \frac{3}{4} = \frac{6}{20}$ b) $\frac{2}{5} \div \frac{1}{3} = \frac{2}{5} \times 3 = \frac{2}{5} \times \frac{3}{1} = \frac{6}{5}$

④ Reduce a fraction.

1. Divide the numerator and denominator by a number that divides evenly into both.

2. Repeat step 1 until no number divides evenly into both the numerator and the denominator.

④ Reduce $\frac{140}{350}$.

1. 10 divides evenly into 140 and into 350. $140 \div 10 = 14$, and $350 \div 10 = 35$. $\frac{140}{350} = \frac{14}{35}$

2. 7 divides evenly into 14 and into 35. $14 \div 7 = 2$, and $35 \div 7 = 5$. $\frac{14}{35} = \frac{2}{5}$

It also would have worked to divide 140 and 350 both by 70 in step 1.

⑤ Add or subtract fractions with the same denominator.

1. Add or subtract the numerators.

2. Keep the same denominator.

⑤ $\frac{12}{20} - \frac{5}{20} = \frac{7}{20}$

⑥ Add or subtract fractions with different denominators.

1. Multiply the first fraction by $\frac{b}{b}$, where b is the denominator of the second fraction.

2. Multiply the second fraction by $\frac{a}{a}$, where a is the denominator of the first fraction.

3. Add or subtract (see ⑤).

⑥ a) $\frac{3}{5} - \frac{1}{4}$

1. $\frac{3}{5} \times \frac{4}{4} = \frac{12}{20}$

2. $\frac{1}{4} \times \frac{5}{5} = \frac{5}{20}$

3. $\frac{12}{20} - \frac{5}{20} = \frac{7}{20}$

$x\%$ means $\frac{x}{100}$. 100% is the same as $\frac{100}{100}$ or 1.

⑦ Convert a percentage to a decimal or fraction.

1. To convert to a fraction, multiply by $\frac{1}{100\%}$, canceling the % symbols. If there is a decimal in the numerator, multiply the fraction by $\frac{10}{10}$ until there is not.
2. To convert to a decimal, move the decimal point two places to the left, and remove the % symbol.

⑦ a) 9%

$$1. \quad 9\% \times \frac{1}{100\%} = \frac{9\%}{100\%} = \frac{9}{100}$$

$$2. \quad 9\% = .09$$

b) .09%

$$.09\% \times \frac{1}{100\%} = \frac{.09\%}{100\%} = \frac{.09}{100}$$

$$\frac{.09}{100} \times \frac{10}{10} = \frac{.9}{1000}$$

$$\frac{.9}{1000} \times \frac{10}{10} = \frac{9}{10,000}$$

$$.09\% = .0009$$

1-B Expressions

The IDENTITY Element of an operation is the value that has no effect under that operation.

For addition and subtraction, the identity element is 0. For multiplication and division, the identity element is 1.

The identity element can also be thought of as the value that remains when after everything is removed, such as $3 - 3 = 0$ or $3 \div 3 = 1$.

A TERM is the product of any number of constants, variables, and function values. For example, 5, $5x^3$, and $5x^3yz^2 \sin x$ are all terms.

The COEFFICIENT of a term is the factor of the term that does not vary, and is usually written at the beginning of the term. The coefficient of each the terms above is 5.

An Algebraic EXPRESSION is one or more terms added together. For example, 5, $5 + 5x^3$, and $5 + 5x^3 - 5x^3yz^2 \sin x$ are all expressions.

① Identify the terms of an expression and their coefficients.

1. The terms are the parts of the equation that equal the whole expression when added together. Terms can be negative, since subtracting is adding a negative.
2. The coefficient of each term is the number that the rest of the term is multiplied by, such as 5 or -5 in the examples above.
 - If no such number is written, then the coefficient is 1 (or -1).
 - If the term is a fraction, the coefficient is a fraction.

① Identify the terms and coefficients of the expression $5x^6 + x^3 + \frac{8x}{7} - 9 - 10\sqrt{x - \cos 3x}$.

1. $5x^6$, x^3 , $\frac{8x}{7}$, -9, and $-10\sqrt{x - \cos 3x}$ are the terms of the expression, because when added the result is the whole expression $5x^6 + x^3 + \frac{8x}{7} - 9 - 10\sqrt{x - \cos 3x}$.

Note that -10, x , $3x$, and $-\cos 3x$ are not terms of this expression.

2. The coefficients are 5, 1, $\frac{8}{7}$, -9, and -10.

An ARGUMENT of a Function is an expression input into the function.

② Identify the argument of a function.

1. Identify the expression being acted upon by the function.

② Identify all arguments in the expression $5x^2 - \frac{8x}{7} + 9 + 10\sqrt{x - \cos 3x}$.

$3x$ is the argument of the cosine function.

$x - \cos 3x$ is the argument of the square root function.

A function and its argument is a single term. An operation on such a term affects the term as a whole, not the argument separately.

③ Multiply a rational expression by an integer.

1. Multiply each term in the numerator by the integer, making sure to avoid the following:

- multiplying a term more than once
- multiplying an argument
- multiplying a term in the denominator

③ Identify the error, if any, in each of the following attempts to multiply $\frac{11(4x) - \sqrt{10x}}{5}$ by 2.

a) $\frac{22(8x) - 2\sqrt{10x}}{5}$ The term $11(3x)$ should be multiplied by 2 one time, but it was multiplied by 2 once on the 11 and again on the $3x$.

b) $\frac{22(4x) - \sqrt{20x}}{5}$ $10x$ is an argument and should not be multiplied.

c) $\frac{22(4x) - 2\sqrt{10x}}{10}$ The denominator should not be multiplied.

d) $\frac{22(4x) - 2\sqrt{10x}}{5}$ This is correct, because each term in the numerator was multiplied by 2 one time.

④ Simplify a fraction with multiple terms in the numerator or denominator.

1. Identify the terms of the numerator and the denominator.

2. Identify a factor that divides evenly into all of the terms.

3. Rewrite the fraction with the factor divided out of each term. If it is variable, specify that it cannot equal zero.

④ $\frac{6x^2 - 9x\sqrt{30x}}{6x^2 - 9x}$

1. The terms are $6x^2$ and $-9x\sqrt{30x}$ in the numerator, and $6x^2$ and $-9x$ in the denominator. $30x$ is an argument of the square root function, not a term of the numerator.

2. $6x^2 \div 3x = 2x$

$-9x\sqrt{30x} \div 3x = -3\sqrt{30x}$ (not $-3\sqrt{10}$)

$6x^2 \div 3x = 2x$

$-9x \div 3x = -3$

3. $\frac{2x - 3\sqrt{30x}}{2x - 3}$, $x \neq 0$ (We divided by $3x$, so our answer doesn't work if $3x$ is zero.)

Note that $6x^2$ in the numerator and $6x^2$ in the denominator do not cancel each other out because there are other terms.

A **MONOMIAL** is a nonzero term with a whole number exponent. The exponent is the **DEGREE**.

<u>Degree</u>	<u>Name</u>	<u>Example</u>
0	CONSTANT	5 (that is, $5x^0$)
1	LINEAR	$5x$ (that is, $5x^1$)
2	QUADRATIC	$5x^2$
3	CUBIC	$5x^3$
n	n^{th} degree	$5x^{12}$

A **POLYNOMIAL** is an expression of the sum of one or more monomials, after like terms are combined. The degree of a polynomial in one variable is the highest degree of its terms.

A **BINOMIAL** is a polynomial with two terms.

A **TRINOMIAL** is a polynomial with three terms.

A polynomial in one variable with its terms written in order from highest degree to lowest is in **STANDARD Form**.

The **LEADING Coefficient** of a Polynomial is the coefficient of the term with the highest degree.

⑤ **Classify a polynomial in one variable.**

1. A single term is a monomial, two terms is a binomial, and three terms is a trinomial.

2. Consider the term with the highest exponent:

If there is no variable at all (that is, there is an unwritten variable with an exponent of 0), the polynomial is degree 0, which is constant.

If there is a variable but no exponent (that is, the variable has an unwritten exponent of 1), the polynomial is degree 1, which is linear.

If the variable is to the second power, the polynomial is quadratic.

If the variable is to the third power, the polynomial is cubic.

If the variable is to the fourth power or higher, look up what it is called, or simply refer to the polynomial as “ n^{th} degree”, where n is the exponent.

⑥ Write the following polynomials in standard form, classify them, and identify the leading coefficient.

	a) $x + 4x^3$	b) $-15x$	c) $8x^2 - 2x^9 + 3$	d) $2 - \frac{7x^3}{5} + 6x^2 + x$
standard form:	$4x^3 + x$	$-15x$	$-2x^9 + 8x^2 + 3$	$-\frac{7x^3}{5} + 6x^2 + x + 2$
classification:	cubic binomial	linear monomial	9 th degree trinomial	cubic polynomial
leading coefficient:	4	-15	-2	$-\frac{7}{5}$

SCIENTIFIC NOTATION is $a \times 10^b$, where $1 \leq a < 10$ and b is an integer.

Many calculators use the notation $a\text{E}b$ instead of $a \times 10^b$ to display scientific notation. Do not write numbers in this notation.

⑥ Convert calculator notation to scientific notation and to standard notation.

1. To convert $a\text{E}b$ to scientific notation, change “E” into “ $\times 10$ ”, and make b an exponent.

2. To convert $a \times 10^b$ to standard notation, move the decimal point right b spaces (which will be left if b is negative). Fill in 0's as needed.

⑥ a) $2.57\text{E}3$ b) $2.57\text{E}-3$

1. 2.57×10^3 2.57×10^{-3}

2. 2570 .00257

⑦ Convert standard notation to scientific notation.

1. Move the decimal place left so that it follows the first nonzero digit, and count how many spaces b it was moved left.

2. Drop any zeros on the end, unless you know they were actually measured, not rounded. If the number is a decimal, no zeros were rounded.

3. Multiply this number by 10^b .

⑦ a) 2700 b) 2700 exactly c) .002700

1. 2.700 ($b = 3$) 2.700 ($b = 3$) 2.700 ($b = -3$)

2. 2.7 2.700 2.700

3. 2.7×10^3 2.700×10^3 2.700×10^{-3}

1-C Solving Equations

When evaluating an expression with no parentheses, exponents are done first, followed by multiplication and division, and then addition and subtraction. When evaluating an expression with parentheses in it, each expression within parentheses is evaluated on its own first, following the order above, before the overall expression is evaluated.

Numerators and denominators always have parentheses around them, even if they are not written. For example, $\frac{1+3}{3+5}$ is actually $\frac{(1+3)}{(3+5)}$. These parentheses only have to be shown if the numerator is not written above the denominator. For example, on a calculator, $\frac{1+3}{3+5}$ can be typed as $(1 + 3) / (3 + 5)$, but not as $1 + 3 / 3 + 5$.

Negative signs mean “multiply by negative 1.” Since they represent multiplication, they are ignored until after exponents are calculated.

① Apply order of operations.

1. Apply the steps below to everything within parentheses, including hidden parentheses.
2. Calculate exponents.
3. Calculate multiplication and division, including multiplying by -1 for negative signs.
4. Calculate addition and subtraction.

① a) $4 + 5 \times 2$

$$4 + 10$$

$$14$$

b) $4 + 5(2)^3$

$$4 + 5(8)$$

$$4 + 40$$

$$44$$

c) -2^4

$$-(16)$$

$$-16$$

d) $(-2)^4$

$$16$$

e) $\frac{5+6}{5-3}$

$$\frac{11}{2}$$

The INVERSE of an operation is the operation that cancels the original operation.

② Identify the inverse of an operation by definition.

1. The inverse of addition is subtraction.
2. The inverse of multiplication is division.
3. The inverse of a power is a root.

② Identify the inverse of each operation in the equation $175x + 900 = 1600$.

The inverse of adding 900 is **subtracting 900**.

The inverse of multiplying by 175 is **dividing by 175**.

An Algebraic EQUATION is one expression set equal to another.

An equation is solved by applying one or more inverse operations to each side of the equation. These are done in reverse order of operations.

③ Solve an equation.

1. If there are parentheses and it is easy to eliminate them (such as by distributing), do so.
 1. Identify the variable you are solving for, or the set of parentheses that includes this variable.
 2. Do each of the following, in order, as needed to isolate the variable or parentheses.
 - Subtract a value or expression from each side to cancel addition, or vice versa.
 - Divide by a value or expression on each side to cancel multiplication, or vice versa.
 - Take the root of each side of the equation to cancel a power, or vice versa.
 3. If one side is now entirely in parentheses, remove the parentheses and redo steps 1-3.

③ a) $3x + 1 = 13$
 $3x = 12$
 $x = 4$

b) $3x^2 + 1 = 13$
 $3x^2 = 12$
 $x^2 = 4$
 $x = \pm 2$

c) $\frac{3}{5}(3x + 1) + 1 = 13$
 $\frac{3}{5}(3x + 1) = 12$
 $\frac{5}{3}(\frac{3}{5}(3x + 1)) = \frac{5}{3}(\frac{12}{1})$
 $(3x + 1) = 20$
 $3x + 1 = 20$
 $3x = 19$
 $x = \frac{19}{3}$

d) $5(3x + 1) + 1 = 4x$
 $15x + 5 + 1 = 4x$
 $6 = 4x - 15x$
 $6 = -11x$
 $x = -\frac{6}{11}$

It is not incorrect to write basic algebraic operations like subtraction or division, such as subtracting 900 or dividing by 175 in the problem below, but there is no need, and it is never done in advanced math. If you do choose to show these, they should be shown as part of each expression, not below them, next to them, or part of only one of them.

Do not use the symbol “ \times ” for multiplication in an equation in which x is a variable.

If t is a variable, make sure it does not look like “+”.

④ Show proper notation in solving an equation.

1. If it is from a word problem, make sure you know what the variable represents.
2. Apply the inverse operations to each side of the equation, not to the equation as a whole or to only one side. Be sure each symbol is written in an appropriate place. Show every step except those that are both simple enough to recognize easily and not new to the current chapter.
3. Write the new equation. Be sure the expressions on either side of the equals sign are equivalent.
4. Repeat steps 2 and 3 as needed.

④ Sam has \$900 this year and plans to save \$175 per year. Solve the equation $175x + 900 = 1600$ to determine when he will have \$1600.

Incorrect

$$175x + 900 = 1600 - 900$$

$$\div 175 (175x = 700)$$

$$x = 700 \div 175 = 4 + 2020 = 2023$$

Reason

The subtraction has to be done on both sides.

An equals sign cannot be divided,
and the \div symbol can't be before the dividend.

$$700 \div 175 \neq 4 + 2020$$

x is years after 2020, not the current year

Correct

x = years after 2020

$$175x + 900 - 900 = 1600 - 900$$

$$175x = 700$$

$$175x \div 175 = 700 \div 175$$

$$x = 4$$

He will have \$1600 in the year $4 + 2020 = 2024$.

④ $\frac{1}{2}x = 5$

There is no work needed to be shown for this problem. Simply writing $x = 10$ makes it clear that both sides were multiplied by 2.

correct but unnecessary: $2(\frac{1}{2}x) = 2(5)$

incorrect: $2(\frac{1}{2}x = 5)$

incorrect: $\times 2 \frac{1}{2}x = 5 \times 2$

incorrect: $\frac{1}{2}x = 5$
 $\times 2 \quad \times 2$

In general, simplifying means writing with smaller numbers if possible, such as by reducing or combining like terms.

Rounding an answer makes it approximate instead of exact. Do not round when asked to simplify.

⑤ Express an answer.

1. If the instructions are to round, then round to the indicated place, or choose an appropriate place to round to if none is specified.
2. If the instructions are to simplify, then reduce fractions, simplify square roots, and combine like terms.
3. If the instructions do not specify whether to round or simplify:
 - If the answer is a number or algebraic expression, do not round unless you have a reason to do so.
 - If the answer is a quantity (that is, it has units, such as *centimeters* or *years*), round unless you have a reason not to do so.

⑤ Give an appropriate answer for each question.

a) Solve $7x = 9$. $x = \frac{9}{7}$

b) Solve $7x = 9$. Round your answer to the nearest tenth. $x = 1.3$

c) If a pack of seven pens costs \$9, what is the price per pen? Each pen costs \$1.29.

1-D Properties of Exponents

The rules below are valid in almost all contexts, including any time a and b are positive or x and y are integers.

Property	Rule	Example
Power of a Product	$(ab)^x = a^x b^x$	$(2x)^3 = 2^3 x^3 = 8x^3$
Power of a Quotient	$\left(\frac{a}{b}\right)^x = \frac{a^x}{b^x}$	$\left(\frac{x}{2}\right)^3 = \frac{x^3}{2^3} = \frac{x^3}{8}$
Power of a Power	$(b^x)^y = b^{xy}$	$(x^5)^3 = x^{15}$
Product of Powers	$b^x b^y = b^{x+y}$	$x^5 x^3 = x^8$
Quotient of Powers	$\frac{b^x}{b^y} = b^{x-y}$	$\frac{x^5}{x^3} = x^2$
Zero Exponent	$b^0 = 1$	$2^0 = 1$
Negative Exponent	$b^{-x} = \frac{1}{b^x}$	$2^{-3} = \frac{1}{2^3} = \frac{1}{8}$

1 Simplify an expression using properties of exponents.

- Use the properties above as needed.
- Reduce fractions if needed.

1 Simplify.

a) $ab^3(a^2b)^5$

$$ab^3(a^{10}b^5)$$

$$a^{11}b^8$$

b) $\left(\frac{5a}{2}\right)^3$

$$\frac{125a^3}{8}$$

c) $\left(\frac{5a}{2}\right)^{-3}$

$$\left(\frac{2}{5a}\right)^3$$

$$\frac{8}{125a^3}$$

d) $10\left(\frac{5a}{2}\right)^{-3}$

$$10\left(\frac{2}{5a}\right)^3$$

$$\frac{10(8)}{125a^3}$$

$$\frac{80}{125a^3}$$

$$\frac{16}{25a^3}$$

e) $\frac{8a^{-3}}{6a^4}$

$$\frac{4}{3a^7}$$

f) $\frac{6 \times 10^{-12}}{2 \times 10^{20}}$

$$3 \times 10^{-32}$$

1 Simplify $\frac{(2a^4b)^3 b^6}{a^{12} b c^2}$, and write it without a fraction. State each property of exponents used.

$$(2a^4b)^3 = 2^3(a^4)^3 b^3$$

Power of a Product

$$\frac{8(a^4)^3 b^3 b^6}{a^{12} b c^2}$$

$$(a^4)^3 = a^{12}$$

Power of a Power

$$\frac{8a^{12} b^3 b^6}{a^{12} b c^2}$$

$$b^3 b^6 = b^9$$

Product of Powers

$$\frac{8a^{12} b^9}{a^{12} b c^2}$$

$$\frac{a^{12} b^9}{a^{12} b} = a^0 b^8$$

Quotient of Powers

$$\frac{8a^0 b^8}{c^2}$$

$$a^0 = 1$$

Zero Exponent

$$\frac{8(1)b^8}{c^2}$$

$$\frac{1}{c^2} = c^{-2}$$

Negative Exponent

$$8b^8 c^{-2}$$

1-E Addition, Subtraction, and Multiplication of Polynomials**① Add or subtract polynomials.**

1. Distribute any coefficient, including negatives.

2. Combine like terms.

① $5(4x^2 + 9x - 3) - (11x - 4)$

1. $(20x^2 + 45x - 15) + (-11x + 4)$

2. $20x^2 + 34x - 11$

② Multiply two polynomials.

1. Multiply each term in the first polynomial by each term in the second polynomial.

2. Simplify.

3. Combine like terms.

② $(4x^2 - 3x)(x + 5)$

1. $(4x^2 \cdot x) + (4x^2 \cdot 5) + (-3x \cdot x) + (-3x \cdot 5)$

2. $4x^3 + 20x^2 - 3x^2 - 15x$

3. $4x^3 + 17x^2 - 15x$

③ Multiply more than two polynomials.

1. Multiply two of the polynomials together (see ②).

2. Multiply the result by the next polynomial.

3. Combine like terms.

③ $(x + 2)(x + 5)(x - 10)$

1. $(x^2 + 7x + 10)(x - 10)$

2. $(x^3 + 7x^2 + 10x) + (-10x^2 - 70x - 100)$

3. $x^3 - 3x^2 - 60x - 100$

The CONJUGATE of a Binomial is the original binomial except with the sign of one of the terms switched: The conjugate of $a + b$ is $a - b$.

There is a special pattern for multiplying a binomial by itself and for multiplying by its conjugate.

Binomial times itself: $(a + b)^2 = a^2 + 2ab + b^2$

Binomial times its conjugate: $(a + b)(a - b) = a^2 - b^2$

④ Multiply a binomial by its conjugate.

1. Square each term of the binomial.
2. Subtract the second square from the first.

④ $(3x - 10)(3x + 10)$

1. $a^2 = (3x)^2 = 9x^2$

$b^2 = 10^2 = 100$

2. $(3x - 10)(3x + 10) = 9x^2 - 100$

⑤ Square a binomial.

1. Square each term of the binomial.
2. Multiply the two terms together, and double this product.
3. Add the three results of the steps above.

⑤ $(3x - 10)^2$

1. $a^2 = (3x)^2 = 9x^2$

$b^2 = (-10)^2 = 100$

2. $2ab = 2(3x)(-10) = -60x$

3. $(3x - 10)^2 = 9x^2 - 60x + 100$